

Improving shower energy resolution for the MINOS charged-current analysis

Fermilab

Christopher Backhouse


University of Oxford

December 13, 2010


Introduction

- ▶ Focusing on disappearance measurements
- ▶ Neutrino oscillations
- ▶ The MINOS experiment
- ▶ The ν_μ analysis
 - ▶ Hadronic energy estimation
- ▶ The $\bar{\nu}_\mu$ analysis
- ▶ Future

Neutrino oscillations - history



- ▶ Atmospheric neutrinos created by cosmic rays striking the atmosphere
- ▶ Super-Kamiokande sees deficit versus expected rate of ν_μ
- ▶ Angle and energy dependent



Neutrino oscillations - theory

$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad i = 1, 2, 3 \quad \alpha = e, \mu, \tau$$

- ▶ PMNS matrix. \sim CKM matrix
for neutrinos

$$P_{\alpha\beta} = \left| \sum_i U_{\alpha i}^* e^{-im_i^2 L/2E} U_{\beta i} \right|^2$$



Neutrino oscillations - theory

$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad i = 1, 2, 3 \quad \alpha = e, \mu, \tau$$

- ▶ PMNS matrix. \sim CKM matrix for neutrinos

$$P_{\alpha\beta} = \left| \sum_i U_{\alpha i}^* e^{-im_i^2 L/2E} U_{\beta i} \right|^2$$

$$U = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$



Neutrino oscillations - theory


$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad i = 1, 2, 3 \quad \alpha = e, \mu, \tau$$

- PMNS matrix. \sim CKM matrix for neutrinos

$$P_{\alpha\beta} = \left| \sum_i U_{\alpha i}^* e^{-im_i^2 L/2E} U_{\beta i} \right|^2$$

$$U = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$P_{\mu\mu} = 1 - \sin^2 2\theta \sin^2 \left(\frac{|\Delta m^2| L}{4E} \right)$$



Neutrino oscillations - theory



$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad i = 1, 2, 3 \quad \alpha = e, \mu, \tau$$

- PMNS matrix. \sim CKM matrix for neutrinos

$$P_{\alpha\beta} = \left| \sum_i U_{\alpha i}^* e^{-im_i^2 L/2E} U_{\beta i} \right|^2$$

$$U = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

$$P_{\mu\mu} = 1 - \sin^2 2\theta \sin^2 \left(\frac{|\Delta m^2| L}{4E} \right)$$



Neutrino oscillations - theory

$$|\nu_\alpha\rangle = \sum_i U_{\alpha i}^* |\nu_i\rangle \quad i = 1, 2, 3 \quad \alpha = e, \mu, \tau$$

► PMNS matrix. \sim CKM matrix for neutrinos

$$P_{\alpha\beta} = \left| \sum_i U_{\alpha i}^* e^{-im_i^2 L/2E} U_{\beta i} \right|^2$$

$$U = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$



$$P_{\mu\mu} = 1 - \sin^2 2\theta \sin^2 \left(\frac{|\Delta m^2| L}{4E} \right)$$

$$\text{NB : } P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) = \mathcal{CP}[P(\nu_\alpha \rightarrow \nu_\beta)]$$

$$P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta) = \mathcal{CPT}[P(\nu_\beta \rightarrow \nu_\alpha)]$$

$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu) = \mathcal{CPT}[P(\nu_\mu \rightarrow \nu_\mu)]$$

C. Backhouse (Oxford)



MINOS

December 13, 2010

Main Injector Neutrino Oscillation Search



- ▶ Long-baseline neutrino oscillation experiment

Main Injector Neutrino Oscillation Search



- ▶ Long-baseline neutrino oscillation experiment
- ▶ Muon neutrino beam from NuMI
- ▶ Near detector on-site at Fermilab
- ▶ Far detector 735 km away in Northern Minnesota

Main Injector Neutrino Oscillation Search



- ▶ Long-baseline neutrino oscillation experiment
- ▶ Muon neutrino beam from NuMI
- ▶ Near detector on-site at Fermilab
- ▶ Far detector 735 km away in Northern Minnesota
- ▶ $L/E \sim 500\text{km/GeV}$
- ▶ Measure ν_μ disappearance


Components of a long-baseline neutrino experiment

- ▶ A neutrino beam
- ▶ A detector technology
 - ▶ Neutrino cross-sections are low
 - ▶ Low cost/kg
- ▶ Two detectors
 - ▶ Near detector to measure beam
 - ▶ Far detector to detect oscillations
 - ▶ Systematics common between the detectors cancel

The NuMI beam




The NuMI beam




- ▶ 120 GeV protons from the Main Injector
- ▶ Strike graphite target
- ▶ Produce hadrons. Primarily π^\pm and K^\pm


The NuMI beam



- ▶ 120 GeV protons from the Main Injector
- ▶ Strike graphite target
- ▶ Produce hadrons. Primarily π^\pm and K^\pm
- ▶ Focused by two magnetic horns
- ▶ Allow us to select charge sign for a neutrino or antineutrino beam



The NuMI beam




- ▶ 120 GeV protons from the Main Injector
- ▶ Strike graphite target
- ▶ Produce hadrons. Primarily π^\pm and K^\pm
- ▶ Focused by two magnetic horns
- ▶ Allow us to select charge sign for a neutrino or antineutrino beam
- ▶ 675 m decay-pipe: $\pi^+ \rightarrow \mu^+ + \nu_\mu$
- ▶ Muons absorbed by rock




The NuMI beam

Total NuMI protons to 00:00 Monday 31 May 2010




- ▶ Taking data since 2005
- ▶ Divided into 3 neutrino run periods, and one antineutrino
- ▶ ν_μ analysis uses an exposure of 7.2×10^{20} protons-on-target
- ▶ $\bar{\nu}_\mu$ analysis uses 1.71×10^{20} POT

Detecting charged particles




- ▶ 4.1 cm × 1 cm plastic scintillator strips
- ▶ TiO₂ coating and embedded wavelength-shifting fibre
- ▶ Scintillation light detected by multi-anode PMTs

Detector construction



U V U V U V U V



- ▶ Magnetized tracking calorimeters
- ▶ Scintillator modules mounted on inch-thick steel planes
- ▶ Alternate scintillator planes at right angles
- ▶ Allows 3D event construction
- ▶ $\mathcal{O}(1\text{T})$ magnetic field → muon charge sign

Near detector



- ▶ 1 kt, $3.8 \text{ m} \times 4.5 \text{ m} \times 16.7 \text{ m}$
- ▶ 1 km from source
- ▶ High event rate: $\sim 10^5$ neutrino interactions per day
- ▶ Measure beam before oscillations


Far detector




- ▶ 5.4 kt, $8\text{ m} \times 8\text{ m} \times 30\text{ m}$
- ▶ 735 km from source, 700 m underground
- ▶ A few neutrino interactions per day
- ▶ Compare spectrum to prediction

Event classes

Charged current




Neutral current




- ▶ Hadronic shower
- ▶ Long muon track
- ▶ Hadronic shower
- ▶ No muon track (possible reconstruction errors)
- ▶ Some energy carried away by outgoing neutrino

Event classes

Charged current




Neutral current




- ▶ Hadronic shower
- ▶ Long muon track
- ▶ Hadronic shower
- ▶ No muon track (possible reconstruction errors)
- ▶ Some energy carried away by outgoing neutrino

Event classes

Charged current




Neutral current




- ▶ Hadronic shower
- ▶ Long muon track
- ▶ Hadronic shower
- ▶ No muon track (possible reconstruction errors)
- ▶ Some energy carried away by outgoing neutrino

Event classes

Charged current



Neutral current



- ▶ High energy muons are easy to pick out
- ▶ Low energy or high-y event → short muon track
- ▶ Likely buried in shower
- ▶ To pick out CC or NC events need to determine nature of reconstructed tracks

Charged current analysis


- ▶ Neutral current events are insensitive to $\nu_\mu \leftrightarrow \nu_\tau$ oscillations
- ▶ Try to get a pure sample of charged current events
- ▶ Look for an energy-dependent deficit

Components of an analysis



- ▶ Select the events you want
- ▶ “Extrapolate” ND spectrum to FD prediction
- ▶ Compare FD data to prediction

- ▶ Plus some improvements since previous analysis




Charged current selection variables







Charged current selection variables



Charged current selection variables




Charged current selection variables





Charged current selector

- ▶ Combine information using a **k-Nearest-Neighbour** algorithm
- ▶ For each event to be classified: find 80 closest Monte Carlo events
- ▶ PID value is the fraction of these neighbours that are charged current



Charged current selector




- ▶ Cut placed at 0.25 - plus another kNN optimized for low energies
- ▶ Above 1 GeV achieve high efficiency with low contamination

Energy estimation

- ▶ $E_{\text{tot}} = E_{\text{trk}} + E_{\text{shw}}$
- ▶ Energy of muon tracks estimated from range
 - ▶ Or curvature if the track exits the detector
- ▶ Previously shower energy estimated calorimetrically
- ▶ New for this analysis: kNN shower energy estimator

Shower energy motivation


MINOS Preliminary



- ▶ Better energy resolution → better resolved oscillation dip → more precise measurement of oscillation parameters

Shower energy motivation

MINOS Preliminary




- ▶ Better energy resolution → better resolved oscillation dip → more precise measurement of oscillation parameters
- ▶ Good gains to be had with modest improvements

Shower energy estimation

- ▶ Track resolution is good.
Shower resolution
dominates →
concentrate on that
- ▶ Use kNN technique
similar to the CC PID

$$\Delta s^2 = \sum_{i=1}^D \frac{(y_i - x_i)^2}{\sigma_i^2}$$

$$E^{\text{reco}} = \frac{1}{k} \sum_{i=1}^k E_i^{\text{true}}$$




Shower energy estimator construction

- ▶ Adjustable parameters are
 - ▶ The input variables (and dimensionality)
 - ▶ The number of neighbours k
- ▶ Figure of merit is Δm^2 errors at maximal mixing
- ▶ Search over possible variable combinations
- ▶ Best kNN has three input variables
 - ▶ Number of planes in the shower
 - ▶ Sum of calorimetric energies of first two showers
 - ▶ Total shower energy within 1 m of the track vertex

Shower energy estimator construction


MINOS Preliminary



- ▶ Search over number of MC neighbours
- ▶ Use 400 neighbours

Improvement


MINOS Preliminary



- ▶ Large improvement at low shower energies


Improvement

MINOS Preliminary



Improvement


MINOS Preliminary



- ▶ Translates to a sensitivity improvement
- ▶ Equivalent to taking 10^{20} POT additional data

Energy corrections


MINOS Preliminary



- ▶ Raw kNN output displays an energy bias
- ▶ Events attracted towards the beam peak in the training set


Energy corrections

MINOS Preliminary



Energy corrections


MINOS Preliminary



- ▶ Fit polynomial function to the bias – divide out

Energy corrections


MINOS Preliminary



- ▶ Fit polynomial function to the bias – divide out
- ▶ Iterate this process → majority of bias corrected

Effect of energy corrections


MINOS Preliminary



- ▶ Energy bias hurts energy resolution because of combination with (unbiased) track energy

Effect of energy corrections


MINOS Preliminary




- ▶ Energy bias hurts energy resolution because of combination with (unbiased) track energy
- ▶ Small improvement from correcting bias

Systematic error on shower energy


- ▶ Contributions to systematic uncertainty from:
 - ▶ Hadronic mismodelling, intranuclear rescattering
 - ▶ Calibration of the detectors
 - ▶ Others...
- ▶ Study hadronic modelling with special MC samples with modified generator



Full set of systematics





Energy calibration



- ▶ Adjust energy variables of input events by $\pm 5.7\%$
- ▶ Leave training events unchanged
- ▶ Do not adjust number of planes
- ▶ Effect smaller at low energies where planes variable more heavily used

Intensity effects and detector gains


- ▶ Possible effect depending on beam intensity
- ▶ Possible effect from detector gains
 - ▶ i.e. overall calibration is right but we lose more or less hits



- ▶ Both effects are small ($< 2\%$)


Final error

- ▶ Sum all contributions in quadrature
- ▶ First crosscheck with calorimetric estimate





Final error

- ▶ Sum all contributions in quadrature
- ▶ First crosscheck with calorimetric estimate






Crosschecks



- ▶ Two contributions also evaluated by reweighting
- ▶ Good agreement


Crosschecks



- ▶ Two contributions also evaluated by reweighting
- ▶ Good agreement
- ▶ Adjust variables to match data
- ▶ Variation is covered by the systematic error band


Resolution binning

- ▶ Another improvement for this analysis
- ▶ Parameterize energy resolution as a function of track and shower energies




$$\frac{\sigma_{\text{trk}}}{E} = \frac{5.1\%}{\sqrt{E}} \oplus 6.9\%$$
$$\frac{\sigma_{\text{shw}}}{E} = \frac{40.4\%}{\sqrt{E}} \oplus 8.6\% \oplus \frac{275\text{MeV}}{E}$$
$$\sigma_{\text{evt}} = \sigma_{\text{trk}} \oplus \sigma_{\text{shw}}$$

Resolution binning




- ▶ Divide events into 5 resolution quantiles
- ▶ Best resolved events give most information about oscillation dip
- ▶ Statistical power of worst events still included

Best 20%




Worst 20%





Extrapolation

- ▶ Use near detector to cancel systematics
- ▶ Measure beam flux, cross-sections and reconstruction effects
- ▶ Need to convert near detector observation into far detector prediction



- ▶ Detectors subtend different solid angles at the decay pipe
- ▶ Neutrinos intersecting the far detector in general come from lower angle pion decays

Extrapolation





$$\text{Flux} \sim \frac{1}{L^2} \left(\frac{1}{1 + \gamma^2 \theta^2} \right)^2$$

$$E_\nu = \frac{0.43 E_\pi}{1 + \gamma^2 \theta^2}$$

- ▶ The same set of parent hadrons give rise to different energy distributions in the two detectors
- ▶ Use Monte Carlo to relate ND neutrinos to their possible parents and thus to the neutrino spectrum at the far detector


Extrapolation



- ▶ Apply a series of corrections to convert observed ND spectrum into ND neutrino flux
- ▶ Use beam matrix to get an FD flux prediction
- ▶ Undo corrections to get an FD spectrum prediction


Oscillation sensitivity

- ▶ Fit is performed using a binned log-likelihood
- ▶ 5 resolution bins fitted separately
- ▶ 90% contour at $\Delta\chi^2 = 4.61$




- ▶ Fit reweighted Monte Carlo sample → sensitivity
- ▶ Add energy estimator and resolution binning sequentially to see effect

Systematic errors





- ▶ Calculate the fit bias introduced by various $\pm 1\sigma$ shift
- ▶ Top four are
 - ▶ Hadronic energy scale
 - ▶ Track energy scale
 - ▶ NC background
 - ▶ Normalization
- ▶ We include these parameters in our fit
- ▶ The result is still statistics-limited

Results - far detector spectrum





- ▶ Observe a deficit of events at low energies

Results - far detector spectrum





- ▶ Observe a deficit of events at low energies
- ▶ Has expected oscillation shape

Results - far detector spectrum





- ▶ Observe a deficit of events at low energies
- ▶ Has expected oscillation shape

Results - far detector spectrum



- ▶ Observe a deficit of events at low energies
- ▶ Has expected oscillation shape
- ▶ Good fit to oscillations


Results - far detector spectrum



- ▶ Observe a deficit of events at low energies
- ▶ Has expected oscillation shape
- ▶ Good fit to oscillations
- ▶ Decay and decoherence don't fit well

Results - contour

MINOS Preliminary



$$|\Delta m^2| = 2.35_{-0.08}^{+0.11} \times 10^{-3} \text{ eV}^2$$


$$\sin^2 2\theta > 0.91 \quad (90\% \text{ C.L.})$$

Pure decay disfavoured $> 6\sigma$


Pure decoherence disfavoured $> 8\sigma$



World's best measurement of Δm_{atm}^2



Introduction to the antineutrino analysis




- ▶ Reverse horn current to select antineutrinos
- ▶ Apply charge-sign cut based on muon curvature
- ▶ Calorimetric shower energy, no resolution binning
- ▶ 90% C.L. determined by Feldman-Cousins procedure
- ▶ Otherwise very similar to neutrino analysis












$$|\Delta \bar{m}^2| = 3.36^{+0.45}_{-0.40} \times 10^{-3} \text{ eV}^2$$



$$\sin^2 2\bar{\theta} = 0.86 \pm 0.11$$







$$|\Delta\bar{m}^2| = 3.36^{+0.45}_{-0.40} \times 10^{-3} \text{ eV}^2$$

$$\sin^2 2\bar{\theta} = 0.86 \pm 0.11$$






$$|\Delta\bar{m}^2| = 3.36^{+0.45}_{-0.40} \times 10^{-3} \text{ eV}^2$$




$$\sin^2 2\bar{\theta} = 0.86 \pm 0.11$$

$$|\Delta m^2| = 2.35^{+0.11}_{-0.08} \times 10^{-3} \text{ eV}^2$$

$$\sin^2 2\theta > 0.91 \quad (90\% \text{ C.L.})$$




Upcoming - rock and antifiducial sample



- ▶ There is a huge amount of mass in the rock around the MINOS cavern
- ▶ And a lot of the detector mass outside the fiducial volume
- ▶ Large number of neutrino interactions
- ▶ Equivalent statistics to fiducial dataset
- ▶ Energy resolution is bad because we only see part of the event
- ▶ Still a valuable sample and sensitivity improvement


Upcoming - more antineutrino running



- ▶ Currently taking additional data in antineutrino mode
- ▶ ~doubling current dataset → significant sensitivity increase

T2K and NO ν A

- ▶ Long baseline neutrino experiments
- ▶ Primarily searching for $\nu_\mu \rightarrow \nu_e$
- ▶ Can also probe ν_μ disappearance



- ▶ T2K - JPARC beam + near detector + SuperKamiokande
- ▶ Started taking data

- ▶ NO ν A - Upgraded NuMI beam
- ▶ Far detector construction started


Conclusion

$$|\Delta m^2| = 2.35^{+0.11}_{-0.08} \times 10^{-3} \text{ eV}^2$$

$$\sin^2 2\theta > 0.91 \quad (90\% \text{ C.L.})$$

$$|\Delta \bar{m}^2| = 3.36^{+0.45}_{-0.40} \times 10^{-3} \text{ eV}^2$$


$$\sin^2 2\bar{\theta} = 0.86 \pm 0.11$$




- ▶ MINOS has the most precise measurements of Δm_{atm}^2 and $\Delta \bar{m}_{\text{atm}}^2$
- ▶ Taking additional antineutrino data now
- ▶ Watch this space...



Backup

ν_e results






MiniBooNE






ND timing cut

MINOS Preliminary




Effect of variables



Effect of variables

MINOS Preliminary



Data driven error

